

Cosmological perturbations in a gravity with quadratic order curvature couplings

Hyerim Noh^(a,c) and Jai-chan Hwang^(b,c)

^(a) Korea Astronomy Observatory, San 36-1, Whaam-dong, Yuseong-gu, Daejeon, Korea

^(b) Department of Astronomy and Atmospheric Sciences, Kyungpook National University, Taegu, Korea

^(c) Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, 85740 Garching bei München, Germany

(February 7, 2008)

We present a set of equations describing the evolution of the scalar-type cosmological perturbation in a gravity with general quadratic order curvature coupling terms. Equations are presented in a gauge ready form, thus are ready to implement various temporal gauge conditions depending on the problems. The Ricci-curvature square term leads to a fourth-order differential equation for describing the spacetime fluctuations in a spatially homogeneous and isotropic cosmological background.

PACS numbers: 04.50.+h, 04.62.+v, 98.80.-k, 98.80.Hw

1. Introduction: In a series of work [1,2] we have been studying the evolution of cosmological perturbations in a gravity with additional curvature-square correction terms

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2} (R + AR^2 + BR^{ab}R_{ab}) + L_m \right], \quad (1)$$

where L_m is the matter part Lagrangian. The gravitational field equation is presented in Eq. (2) of [1]. The one-loop order quantum correction generically leads to curvature-square terms, and due to Gauss-Bonnet theorem the two terms in Eq. (1) completely account such corrections, [3]. In the literature, many works analyzed the roles of R^2 term in the context of cosmological structure formation, [4]. However, we find rarely any work considering the Ricci-curvature square term in the same context. When the curvature-square terms come from quantum correction of Einstein gravity, if R^2 term is important $R^{ab}R_{ab}$ term should be important as well because two terms are of the same order.

As a background world model we consider a spatially flat, homogeneous, and isotropic, Friedmann-Lemaître-Robertson-Walker (FLRW), model. The general perturbations of the FLRW spacetime can be decomposed into three-types based on the tensorial properties of the perturbed variables: these are the scalar-type, (transverse) vector-type, and (transverse-tracefree) tensor-type perturbations. The tensor- and vector-type perturbations correspond to the gravitational wave and the rotational perturbation, respectively. The scalar-type perturbation includes the density condensation and the corresponding longitudinal type velocity perturbation, gravitational potential (curvature) perturbation, etc. Due to the high symmetry in the background model (homogeneity and isotropy) the three-types of perturbations decouple from each other in the linear level and evolve independently. Thus, without losing generality we can consider the three perturbations independently. The gravitational wave and the rotational perturbations were analysed in [1] and [2], respectively. For the gravitational wave we derived a fourth-order differential equation, whereas, the rotational perturbation is simply described by the conservation of angular momentum. In this *Brief Report*, we will present a complete set of equations for the scalar-type perturbation in a gauge-ready form: these are in Eqs. (8-14). We will show that, in general, the Ricci-curvature square term leads to a fourth-order differential equation. Applications to specific cosmological scenarios are left for future work.

2. Scalar-type perturbation: The cosmological spacetime metric with the most general scalar-type perturbation is

$$ds^2 = -a^2(1 + 2\alpha)d\eta^2 - a^2\beta_{,\alpha}d\eta dx^\alpha + a^2[\delta_{\alpha\beta}(1 + 2\varphi) + 2\gamma_{,\alpha\beta}]dx^\alpha dx^\beta, \quad (2)$$

where $a(\eta)$ is a cosmic scale factor. $\alpha(\mathbf{x}, \eta)$, $\beta(\mathbf{x}, \eta)$, $\varphi(\mathbf{x}, \eta)$ and $\gamma(\mathbf{x}, \eta)$ are perturbed order metric variables. Except for two degrees of the gauge redundancy, these variables represent the scalar-type metric perturbations. It is convenient to introduce the following combinations $\chi \equiv a(\beta + a\dot{\gamma})$ and $\kappa \equiv 3(H\alpha - \dot{\varphi}) - (\Delta/a^2)\chi$; Δ is a Laplacian based on $\delta_{\alpha\beta}$, an overdot denotes the time derivative based on t where $dt \equiv a d\eta$, and $H \equiv \dot{a}/a$. α is the perturbed parts of the comoving lapse function. φ , χ , and κ are the perturbed parts of the three-space curvature, the shear, and minus of the perturbed expansion scalar of the normal frame vector field, respectively; see Sec. 2.1.1 in [5]. We decompose $R(\mathbf{x}, t) = \bar{R}(t) + \delta R(\mathbf{x}, t)$; unless necessary, we ignore the overbar which indicates the quantity to the background order. For later use we present:

$$R_0^0 = 3(\dot{H} + H^2) - \left(\dot{\kappa} + 2H\kappa + 3\dot{H}\alpha + \frac{\Delta}{a^2}\alpha \right), \quad R_\alpha^0 = -\frac{2}{3}\frac{1}{a} \left(\kappa + \frac{\Delta}{a^2}\chi \right)_{,\alpha},$$

$$R_\beta^\alpha = \left(\dot{H} + 3H^2 \right) \delta_\beta^\alpha - \frac{1}{3} \left[\dot{\kappa} + 6H\kappa + 3\dot{H}\alpha + \frac{\Delta}{a^2} (3\varphi + \dot{\chi} + H\chi) \right] \delta_\beta^\alpha - \frac{1}{a^2} (\varphi + \alpha - \dot{\chi} - H\chi)^{|\alpha|}_\beta, \quad (3)$$

where $0 = \eta$ and $|$ indicates a covariant derivative based on $\delta_{\alpha\beta}$.

3. Gauge strategy: The gauge transformation properties of the metric variables are presented in Sec. 2.2 of [5]. Under the gauge transformation $\tilde{x}^a = x^a + \xi^a$ with $T \equiv a\xi^0$, we have [see Eq. (17) in [5]]:

$$\tilde{\alpha} = \alpha - \dot{T}, \quad \tilde{\varphi} = \varphi - HT, \quad \tilde{\chi} = \chi - T, \quad \tilde{\kappa} = \kappa + (3\dot{H} + \Delta/a^2)T, \quad \delta\tilde{R} = \delta R - \dot{R}T. \quad (4)$$

The variables α , φ , χ , κ , and δR only depend on the temporal gauge transformation, thus are spatially gauge invariant; for δR we used Eq. (3), see also Eq. (14). We have several choices for the temporal gauge fixing: the synchronous gauge ($\alpha \equiv 0$), the uniform-curvature gauge ($\varphi \equiv 0$), the zero-shear gauge ($\chi \equiv 0$), the uniform-expansion gauge ($\kappa \equiv 0$), and the uniform- R gauge ($\delta R \equiv 0$). Unless we consider perturbations in the energy-momentum tensor we do not have the comoving gauge (or the uniform-field gauge for the scalar field), the uniform-density gauge, etc., which are related to imposing conditions on δT_b^a . Except for the synchronous gauge condition, each of other temporal gauge conditions *completely fixes* the temporal gauge mode. Thus, a variable in such a gauge condition corresponds to a unique gauge invariant combination involving the variable concerned and the variable used in the gauge condition. We proposed the following notation for the gauge invariant variables [5]:

$$\varphi_\chi \equiv \varphi - H\chi \equiv -H\chi_\varphi, \quad \delta R_\chi \equiv \delta R - \dot{R}\chi \equiv -\dot{R}\chi_{\delta R}, \quad \delta R_\varphi \equiv \delta R - \frac{\dot{R}}{H}\varphi \equiv -\frac{\dot{R}}{H}\varphi_{\delta R}, \quad \text{etc.} \quad (5)$$

In this manner we can construct systematically all possible gauge invariant combinations. Since there exist several (in principle, infinitely many) different gauge conditions available for each variable, and correspondingly several gauge invariant combinations, our notation for the gauge invariant variable is convenient in practice.

In the Ricci-curvature square (we call it Ricci-square) gravity we have several additional temporal gauge conditions. From Eqs. (3,4) we have:

$$\delta\tilde{R}_0^0 = \delta R_0^0 - \dot{R}_0^0 T, \quad \delta\tilde{R}_\alpha^0 = \delta R_\alpha^0 - \frac{2}{a}\dot{H}T_{,\alpha}, \quad \delta\tilde{R}_\beta^\alpha = \delta R_\beta^\alpha - \frac{1}{3}\dot{R}_\gamma^\alpha \delta_\beta^\gamma T. \quad (6)$$

Thus, a temporal gauge condition among $\delta R_0^0 = 0$, $\delta R_\gamma^{0\gamma} = 0$, and $\delta R_\gamma^\gamma - \delta R_0^0 = 0$ also completely fix the temporal gauge degree of freedom; but, these are combinations of our fundamental gauge conditions, because these conditions are the same as $\delta R - 4[H\kappa + (\Delta/a^2)\varphi] = 0$, $\kappa + (\Delta/a^2)\chi = 0$, and $H\kappa + (\Delta/a^2)\varphi = 0$, respectively,

In the following section we will present the perturbed set of equations *without imposing* the temporal gauge condition. In this way, the right to impose the temporal gauge condition can be used as an *advantage* for handling problems depending on the mathematical simplification or the physical interpretation we can achieve [6,5]. Often, different gauge conditions suit for different problems and keeping the equations ready for imposing various gauge conditions (thus, in a gauge ready form) is practically convenient.

4. Equations in the gauge ready form: Equations for the background are presented in Eq. (7) of [1]:

$$H^2 + 2(3A + B) \left(2H\ddot{H} - \dot{H}^2 + 6H^2\dot{H} \right) = \frac{1}{3}\mu, \quad \dot{\mu} = -3H(\mu + p), \quad \dot{H} = -\frac{1}{2\mathcal{F}} \left(\ddot{\mathcal{F}} - \dot{\mathcal{F}} + \mu + p \right), \quad (7)$$

where $\mathcal{F} \equiv 1 + 2(A + B/3)R$, $\mu = -\bar{T}_0^0$, and $p = \frac{1}{3}\bar{T}_\gamma^\gamma$; T_{ab} is the energy-momentum tensor of the additional matter part Lagrangian. The third equation follows from the first two equations.

The perturbed set of equations follows from the gravitational field equation in Eq. (2) of [1]. The complete set of equations in the gauge ready form is the following [we introduce $F \equiv 1 + 2AR$, thus $\delta F = 2A\delta R$]:
Definition of κ :

$$\kappa \equiv -3(\dot{\varphi} - H\alpha) - \frac{\Delta}{a^2}\chi. \quad (8)$$

Energy constraint:

$$\delta T_0^0 = 2F \left[\frac{\Delta}{a^2}\varphi + \left(H + \frac{\dot{F}}{2F} \right) \kappa + \frac{3H\dot{F}}{2F}\alpha \right] - 3H\delta\dot{F} + \left[3 \left(\dot{H} + H^2 \right) + \frac{\Delta}{a^2} \right] \delta F$$

$$\begin{aligned}
& -B \left\{ 2 \left(H\kappa + \frac{\Delta}{a^2} \varphi \right)'' + 6H \left(H\kappa + \frac{\Delta}{a^2} \varphi \right)' + 2 \left[2 \left(\dot{H} - 6H^2 \right) - \frac{\Delta}{a^2} \right] \left(H\kappa + \frac{\Delta}{a^2} \varphi \right) + 3H\delta\dot{R} - \left(3H^2 + \frac{\Delta}{a^2} \right) \delta R \right. \\
& \quad \left. + 6H\dot{H}\dot{\alpha} + 6 \left(-H\ddot{H} + 2\dot{H}^2 - 9H^2\dot{H} \right) \alpha - \left(6\dot{H} + 5H^2 \right)' \kappa - \frac{8}{3}H\frac{\Delta}{a^2} \left(\kappa + \frac{\Delta}{a^2} \chi \right) \right\}. \tag{9}
\end{aligned}$$

Momentum constraint:

$$\begin{aligned}
T_\alpha^0 &= \frac{2}{3} \frac{1}{a} \nabla_\alpha \left[\left[-F \left(\kappa + \frac{\Delta}{a^2} \chi + \frac{3\dot{F}}{2F} \alpha \right) + \frac{3}{2} \left(\delta\dot{F} - H\delta F \right) \right] \right. \\
& \quad \left. -B \left\{ - \left(\kappa + \frac{\Delta}{a^2} \chi \right)'' - 3H \left(\kappa + \frac{\Delta}{a^2} \chi \right)' + \left(12H^2 + \frac{\Delta}{a^2} \right) \left(\kappa + \frac{\Delta}{a^2} \chi \right) - \frac{3}{2} \left(\delta\dot{R} - H\delta R \right) \right. \right. \\
& \quad \left. \left. - 3\dot{H}\dot{\alpha} + \frac{3}{2} \left(2\dot{H} + 9H^2 \right)' \alpha + 3\frac{\Delta}{a^2} \left[2H\varphi + \left(\dot{H} - 2H^2 \right) \chi \right] \right\} \right]. \tag{10}
\end{aligned}$$

Tracefree propagation:

$$\begin{aligned}
\delta T_\beta^\alpha - \frac{1}{3} \delta T_\gamma^\gamma \delta_\beta^\alpha &= \frac{1}{a^2} \left(\nabla^\alpha \nabla_\beta - \frac{1}{3} \delta_\beta^\alpha \Delta \right) \left[F \left[\dot{\chi} + \left(H + \frac{\dot{F}}{F} \right) \chi - \varphi - \alpha - \frac{\delta F}{F} \right] \right. \\
& \quad \left. -B \left\{ \left(\dot{\chi} + H\chi - \varphi - \alpha \right)'' - H \left(\dot{\chi} + H\chi - \varphi - \alpha \right)' - \left[8 \left(\dot{H} + H^2 \right) + \frac{\Delta}{a^2} \right] \left(\dot{\chi} + H\chi - \varphi - \alpha \right) \right. \right. \\
& \quad \left. \left. + \delta R - 4\dot{H}\varphi - 6 \left(\dot{H} + H^2 \right)' \chi + \frac{8}{3}H \left(\kappa + \frac{\Delta}{a^2} \chi \right) \right\} \right]. \tag{11}
\end{aligned}$$

Raychaudhuri equation:

$$\begin{aligned}
\delta T_\gamma^\gamma - \delta T_0^0 &= 2F \left\{ \dot{\kappa} + \left(2H + \frac{\dot{F}}{2F} \right) \kappa + \frac{3\dot{F}}{2F} \dot{\alpha} + \left[3\dot{H} + \frac{3}{2F} \left(2\ddot{F} + H\dot{F} \right) + \frac{\Delta}{a^2} \right] \alpha \right\} \\
& \quad - 3\delta\ddot{F} - 3H\delta\dot{F} + \left(6H^2 + \frac{\Delta}{a^2} \right) \delta F \\
& \quad -B \left\{ -4 \left(H\kappa + \frac{\Delta}{a^2} \varphi \right)'' - 12H \left(H\kappa + \frac{\Delta}{a^2} \varphi \right)' - 4 \left[2 \left(\dot{H} - 6H^2 \right) - \frac{\Delta}{a^2} \right] \left(H\kappa + \frac{\Delta}{a^2} \varphi \right) + 2\delta\ddot{R} + 6H^2\delta R \right. \\
& \quad \left. - 28H\dot{H}\kappa - 6 \left(2\dot{H} + 5H^2 \right)' \dot{\alpha} - 12 \left[\left(2\dot{H} + 5H^2 \right)'' + 3H^2\dot{H} \right] \alpha + \frac{16}{3}H\frac{\Delta}{a^2} \left(\kappa + \frac{\Delta}{a^2} \chi \right) \right\}. \tag{12}
\end{aligned}$$

Trace equation:

$$\delta T = -\delta R - 2(3A + B) \left[\delta\ddot{R} + 3H\delta\dot{R} - \frac{\Delta}{a^2} \delta R - \dot{R}(\kappa + \dot{\alpha}) - \left(2\ddot{R} + 3H\dot{R} \right) \alpha \right], \tag{13}$$

and we have

$$\delta R = -2 \left[\dot{\kappa} + 4H\kappa + 3\dot{H}\alpha + \frac{\Delta}{a^2} (2\varphi + \alpha) \right]. \tag{14}$$

We decomposed the equation for δT_β^α into tracefree and trace parts. Equations (8-14) form a redundantly complete set describing the scalar-type perturbation in a gauge ready form; for example, the combination of Eqs. (9,12) leads to Eq. (13). In the limit of R^2 gravity (thus $B = 0$) Eqs. (8-13) correspond to Eqs. (54-58,60) in [7] which also correspond to Eqs. (22-26) in [5]; the names of the equations are based on our convention in [5]. When we analyse we have the right to impose one temporal gauge condition out of several choices displayed in Sec. 3.

5. R^2 gravity: We set $B = 0$. Without additional matter terms ($T_{ab} = 0$), the analysis can be made most conveniently in terms of φ in the uniform- R gauge where we take $\delta R \equiv 0$ (thus $\delta F = 0$) as the gauge condition;

equivalently, we let $\delta R = 0$ and change the other perturbation variables into the gauge invariant forms, for example, $\varphi \rightarrow \varphi_{\delta R}$, etc. Using Eq. (8-10) we express κ and α in terms of φ . From Eq. (13) we can derive

$$\frac{(HF/\dot{F} + 1/2)^2}{a^3 F} \left[\frac{a^3 F}{(HF/\dot{F} + 1/2)^2} \dot{\varphi}_{\delta R} \right] - \frac{\Delta}{a^2} \varphi_{\delta R} = 0. \quad (15)$$

This equation was derived in a more general context in [8,7]; it is valid for a general action with $f(R)$ replacing $R + AR^2$ with $F \equiv \partial f / (\partial R)$. Perturbation analyses in the R^2 gravity were also pursued in [4] in different gauge conditions. Using $\varphi_{\delta R} = -(H/\dot{R})\delta R_\varphi$ in Eq. (5) we can derive the equation for δR_φ . In the large scale limit (thus, ignoring the Laplacian term) we have an integral form *solution valid for the generally time varying background dynamics*

$$\varphi_{\delta R}(\mathbf{x}, t) = -\frac{H}{\dot{R}} \delta R_\varphi(\mathbf{x}, t) = C(\mathbf{x}) + \frac{2}{3} D(\mathbf{x}) \int_0^t \frac{(HF/\dot{F} + 1/2)^2}{a^3 F} dt, \quad (16)$$

where $C(\mathbf{x})$ and $D(\mathbf{x})$ are integration constants indicating the growing and decaying solutions, respectively; the decaying solution is higher order in the large scale expansion, see Eq. (109) in [7]. Thus, the non-transient solution of $\varphi_{\delta R}$ is conserved in the large scale limit, and the generalized nature of the gravity does not affect this result!

6. Discussions: In general we expect the curvature square terms lead to fourth-order gravity theories. It happens that, as we just saw, the R^2 or $f(R)$ gravity leads to a second-order system. This is probably due to the conformal symmetry of the $f(R)$ gravity to the Einstein gravity with a minimally coupled scalar field, [9]. The Ricci-square term does not have such a conformal symmetry, and we expect to have a fourth-order differential equation for the scalar-type perturbation. In general, we do not know which gauge condition will be the most suitable for the problem *a priori*. As a trial attempt we can impose the same gauge condition used in R^2 gravity which is the uniform- R gauge; thus, we set $\delta R \equiv 0$ and $\delta F = 0$. One possible way to derive a fourth-order equation is the following. Using Eqs. (8,13,14) we can express α , $\dot{\alpha}$, and $\kappa + (\Delta/a^2)\chi$ in terms of κ and φ . From a time derivative of Eq. (14), and from Eq. (9) we can derive a coupled set of two second-order differential equations for κ and $H\kappa + (\Delta/a^2)\varphi$. In such a special form, however, it happens that the equations alone cannot be reduced to Eq. (15) in a pure R^2 gravity limit.

Our *main result* is Eqs. (8-14) which describe the evolution of a perturbed flat FLRW background in a gauge ready form; the evolution of background is governed by Eq. (7). The set of equations for the Ricci-square term is a new result, and after imposing a suitable gauge condition it generally leads to a fourth-order differential equation without remaining gauge mode. We mentioned one possible set of a fourth-order differential equation in a previous paragraph as a demonstration. However, the mentioned set may not be the best form possible in the available pool of the variables and the gauge conditions in Sec. 3. Equations (8-14) are arranged in a general gauge-ready form, thus allow an easy implementation of one's favored choice for the variables and the temporal gauge.

The fourth-order terms in Eq. (1) usually appear as the one-loop order quantum corrections. In this regard such terms can be regarded as a transient (because higher-order correction terms will appear soon in higher order quantum corrections) form of medium-energy effective action [3]. Thus, we regard the A and B terms in Eq. (1) as corrections to Einstein action, and in such a case, we should consider the third and fourth order derivatives as corrections to the second order equation based on Einstein gravity. In the case we should accept Eq. (1) as the fundamental theory we are not certain whether the possible four solutions of the expected fourth-order differential equation in a general situation (and also the fourth-order gravitational wave equation derived in [1]) should be interpreted as representing physical degrees of freedom; for example, we should consider the Cauchy problem and the stability of the full theory. We will not address this important question in this *Brief Report*, [10].

We briefly describe how we can connect specific solutions of our formulation with the present day observable quantities. Since we have some conceptual difficulties mentioned in the previous paragraph for fourth-order gravity, let us consider the R^2 gravity as an example. Let us consider a scenario where the higher-order gravity dominates the early evolution stage of our observable patch of the universe, and at some point Einstein gravity takes over the dominance till the present era. If we further assume that the higher-order gravity era provides an accelerated expansion (inflation) stage, the observationally relevant scales may transit from subhorizon to superhorizon scales during the stage. If conditions (e.g., allowing an analytic form mode-function solution of the perturbed equation) are met, we can derive the generated quantum fluctuations (usually, in general scales) based on vacuum expectation value, and as the scale becomes superhorizon it can be interpreted as the classical fluctuations based on the spatial average. As long as the relevant scales remain in the superhorizon stage during the transit epoch of the gravities (from higher-order to Einstein) there exists a *conserved quantity*. φ in the comoving gauge is the conserved quantity in the superhorizon scale independently of the changing gravity (as long as the gravity belongs to a certain class), changing field potential (for the scalar field), and changing equation of state (for the hydrodynamic situation), see [8]. In our case, $\varphi_{\delta R}$ in Eq. (16) is the conserved quantity; the growing solution is conserved as $C(\mathbf{x})$. Since we are considering the linear

perturbation all quantities are related to each other by linear combinations; for the specific relations to the quantities with observational consequences (e.g., temperature anisotropies in the cosmic microwave background radiation, and the Newtonian hydrodynamic variables) see Eq. (23) in [8]. In summary, the curvature fluctuation φ in a certain gauge is conserved and connects perturbations in both eras. In later Einstein era with hydrodynamic contents, the density fluctuations will fall-in to the potential well of, thus co-grow with, the curvature fluctuation. Concrete analyses of the quantum generation and classical evolution processes with the generated observational spectra in various scenarios based on a class of generalized gravity theories were carried out in [11]. Applications of our formulation to specific cosmological models are left for the future work.

JH was supported by the KOSEF, Grant No. 95-0702-04-01-3 and through the SRC program of SNU-CTP. HN was supported by the DFG fellowship (Germany) and the KOSEF (Korea).

-
- [1] H. Noh and J. Hwang, Phys. Rev. D **55**, 5222 (1997).
 - [2] J. Hwang and H. Noh, Phys. Rev. D **57**, 2617 (1998).
 - [3] A. Pais and G. E. Uhlenbeck, Phys. Rev. **79**, 145 (1950); R. Utiyama and B. DeWitt, J. Math. Phys. **3**, 608 (1962); B. S. DeWitt, *Dynamical theory of groups and fields* (New York, Gordon and Breach, 1965); A. D. Sakharov, Sov. Phys. Dokl. **12**, 1040 (1968); G. 't Hooft and M. Veltman, Ann. Inst. Henri Poincaré **XX**, 69 (1974); K. S. Stelle, Gen. Rel. Grav. **9**, 353 (1978).
 - [4] H. Nariai, Prog. Theor. Phys. **49**, 165 (1973); L. A. Kofman and V. F. Mukhanov, JETP Lett. **44**, 619 (1986); V. F. Mukhanov, L. A. Kofman and D. Yu. Pogosyan, Phys. Lett. B **193**, 427 (1987); L. A. Kofman, V. F. Mukhanov and D. Yu. Pogosyan, Sov. Phys. JETP **66**, 433 (1988); V. F. Mukhanov, Phys. Lett. B **218**, 17 (1989); M. B. Baibosunov, V. Ts. Gurovich and U. M. Imanaliev, Sov. Phys. JETP **71**, 636 (1990); J. Hwang, Class. Quantum Grav. **7**, 1613 (1990); Phys. Rev. D **42**, 2601 (1990); Class. Quantum Grav. **8**, 195 (1990); V. F. Mukhanov, H. A. Feldman and R. H. Brandenberger, Phys. Rep. **215**, 203 (1992).
 - [5] J. Hwang, Astrophys. J. **375**, 443 (1991).
 - [6] J. M. Bardeen, in *Particle Physics and Cosmology*, edited by L. Fang and A. Zee (Gordon & Breach, London, 1988), 1p.
 - [7] J. Hwang and H. Noh, Phys. Rev. D **54**, 1460 (1996).
 - [8] J. Hwang, Phys. Rev. D **53**, 762 (1996).
 - [9] R. H. Dicke, Phys. Rev. **125**, 2163 (1962); B. Whitt, Phys. Lett. B **145**, 176 (1984).
 - [10] J. Z. Simon, Phys. Rev. D **41**, 3720 (1990); *ibid.* **43**, 3308 (1991); J. F. Donoghue, *ibid.* **50**, 3874 (1994).
 - [11] J. Hwang, Class. Quantum Grav. **14**, 3327 (1997); J. Hwang and H. Noh, *ibid.* **15**, 1387 (1998).